

Spectral Energy Distributions of Flat Spectrum Radio Quasars observed with *BeppoSAX*.

Tavecchio F.¹, Maraschi L.¹, Ghisellini G.², Celotti A.³, Chiappetti L.⁴, Comastri A.⁵,
Fossati G.⁶, Grandi P.⁷, Pian E.⁸, Tagliaferri G.², Treves A.⁹, Sambruna R.¹⁰

Received _____; accepted _____

¹Osservatorio Astronomico di Brera, via Brera 28, 20121, Milano, Italy.

²Osservatorio Astronomico di Brera, via Bianchi 46, 23807 Merate, Italy

³SISSA/ISAS, via Beirut 2-4, 34014 Trieste, Italy

⁴IFC/CNR, via Bassini 15, 20133, Milano, Italy

⁵Osservatorio Astronomico di Bologna, via Ranzani 1, 40127, Bologna, Italy

⁶CASS, University of California, San Diego, CA 92093-0424

⁷IAS, IAS/CNR, via Fosso del Cavaliere, 00133 Roma, Italy

⁸Osservatorio Astronomico di Trieste, via Tiepolo 11, 24131 Trieste, Italy

⁹Universita' dell'Insubria, via Valleggio 11, 22100, Como, Italy

¹⁰George Mason University, 4400 University Dr., M/S 3F3, Fairfax, VA, 22030-4444

ABSTRACT

We report the *BeppoSAX* observations of 6 Flat Spectrum Radio Quasars. Three of them have a clear detection up to 100 keV with the PDS instrument. For 4 objects the X-ray spectrum is satisfactorily fitted by a power-law continuum with Galactic absorption. 2251+158 show the presence of absorption higher than the galactic value, while the spectrum of the source 0208-512 shows a complex structure, with evidence of absorption at low energy. We construct the Spectral Energy Distributions adding historical data to the broad band X-ray spectra obtained with *BeppoSAX* and reproduce them with a one-zone Synchrotron-Inverse Compton model (including both SSC and External Compton). The implications are briefly discussed.

Subject headings: quasars: general — quasars: individual (0208-512, 0521-365, 1641+399, 2223-052, 2243-123, 2251+158) — X-rays: spectra — radiation mechanisms: non-thermal

1. Introduction

Blazars are the best laboratory to study the physics of relativistic jets. The non-thermal continuum amplified by relativistic beaming (e.g., Urry & Padovani 1995), is a unique tool to probe the physical processes acting in the jet. In the last decade the interest for Blazars has been renewed by the EGRET discovery of 66 Blazars (Hartman et al. 1999) as strong γ -ray emitters. In several cases the γ -ray emission dominates the apparent bolometric luminosity. Moreover the short variability timescales, together with the absence of absorption of high energy photons (through the $\gamma\gamma$ -pair production), directly imply that the γ -ray emission is also relativistically beamed (e.g. Dondi & Ghisellini 1995).

Despite the large variety of classifications, there is growing evidence that Blazars form a single population. Their different spectral properties can be unified within a spectral sequence in which the leading parameter is the total luminosity (Fossati et al. 1998). At the high-luminosity extreme of the sequence we find the Flat Spectrum Radio Quasars, the most luminous blazars, characterized by the presence of bright emission lines in the optical-UV spectrum and, in some cases, strong *UV bumps*, signatures of the underlying accretion process (e.g. Pian et al. 1999 and references therein). Most of the Blazars detected by EGRET (Hartman et al. 1999) belong to this sub-class but not all FSRQs with comparable SEDs have been detected in the γ -ray domain.

In the study of FSRQs, observations in the X-ray band play a fundamental role. The X-ray emission from FSRQ is dominated by the continuum originating from Inverse Compton scattering between relativistic electrons in the jet and the soft photons produced by the disk and/or in the Broad Line Region (Dermer & Schlickeiser 1993; Sikora, Begelman & Rees 1994). The possibility to measure the IC component yields important constraints on the emission models. Moreover the study of the soft X-ray spectrum (below 1 keV) provides insight on the presence of intrinsic absorption, on the possible contribution of the high

energy end of the synchrotron emission or on the minimum energy of emitting particles. The latter point is particularly interesting because the determination of the minimum energy of particles is a fundamental step in the estimate of the total power transported by the jet (e.g. Celotti, Padovani & Ghisellini 1997).

The *BeppoSAX* satellite has the unique capability of covering the wide energy range 0.1 – 200 keV. For this reason it is ideal to study FSRQs. In particular the knowledge of the spectrum in this band, together with the γ -ray spectrum measured by EGRET, provide the most complete spectral information on the high energy component (see e.g., Tavecchio et al. 2000, hereafter Paper I). For these reasons we started an observational program with *BeppoSAX* of a sample of FSRQs, (containing a total of 50 sources) extracted from the 2-Jy sample by Padovani & Urry (1992). Using the flux threshold $F_{1\text{ keV}} > 0.5\mu\text{Jy}$ the number of sources reduces to 19. In Paper I we discussed the case of three FSRQ detected by EGRET, namely 0836+710, 1510-089 and 2230+114. Here we report the analysis of *BeppoSAX* data of 6 other sources, 3 detected by EGRET and three without an EGRET detection. Adding these 6 sources to the 3 FSRQs analyzed in Paper I and other 3 sources (3C279, 0528+134, PKS 0537-441) discussed elsewhere (Maraschi et al. 1998; Ghisellini et al. 1999, Pian et al., in prep) a total of 12 FSRQs (out of 19 with X-ray flux $F_{1\text{ keV}} > 0.5\mu\text{Jy}$) from the 2 Jy sample were observed by *BeppoSAX*. The sources discussed here are listed in Table 1. Partial and preliminary results were reported in Maraschi & Tavecchio (2001)

The paper is organized as follows: in section 2 we report the analysis of *BeppoSAX* observations, in section 3 we discuss the emission models for the SEDs and finally in Sect. 4 we discuss our results.

2. *BeppoSAX* observations and Analysis

The scientific payload of the *BeppoSAX* satellite (see Boella et al. 1997) contains four coaligned Narrow Field Instruments (NFIs) and two Wide Field Cameras. Two of the NFIs use concentrators to focalize X-rays: the Low Energy Concentrator Spectrometer (LECS) has a detector sensitive to soft-medium X-ray photons (0.1-10 keV), while the Medium Concentrator Spectrometer (MECS) can detect photons in the medium energy range 1.3-10 keV. The Phoswich Detector System (PDS), sensitive from 12 up to 200 keV, consisting of four identical units, uses rocking collimators so as to monitor source and background simultaneously with interchangeable units. We will not be concerned here with the fourth NFI, a High Pressure Gas Scintillation Proportional Counter (HPGSPC).

The *BeppoSAX* journal of observations is reported in Table 2, with exposure times and observed count rates. None of the sources showed significant flux variations during the observations. We therefore obtained a cumulative spectrum for each source.

We analyzed the *BeppoSAX* spectral data using the standard software packages XSELECT (v 1.4b) and XSPEC (v 11.0) and the September 97 version of the calibration files released by the *BeppoSAX* Scientific Data Center (SDC). From the event files we extracted the LECS and MECS spectra in circular regions centered around the source with radii of 8' and 4' respectively (see the SAX Analysis Cookbook¹). The PDS spectra extracted with the standard pipeline with the rise-time correction were directly provided by the *BeppoSAX* SDC. We used PDS data rebinned with $S/N > 3$.

For the spectral analysis we considered the LECS data in the restricted energy range 0.1-4 keV, because of known unsolved problems with the response matrix at higher energies. Background spectra extracted from blank field observations at the same position as the

¹ftp://www.sdc.asi.it/pub/sax/doc/software_docs/saxabc_v1.2.ps.gz

source were used. We fitted rebinned LECS, MECS and PDS spectra jointly, allowing for two variable different normalization factors to take into account uncertainties in the intercalibration of different instruments (see SAX Cookbook).

The first model chosen for the fitting procedure was in all cases a single power law plus Galactic absorption. In the following we discuss in detail the results for each source. The results of the spectral fits are summarized in Table 3 and 4.

2.1. 0208-512

The residuals of a simple power-law model to the LECS+MECS data show the presence of a prominent absorption feature, located at ~ 0.6 keV, (see Fig.1) and an excess at about 5 keV. Inspection of the individual MECS2 and MECS3 spectra reveals that the emission feature at 5 keV is present only in the data obtained with the MECS3 indicating an instrumental origin. Only the MECS2 data are used below.

The low energy feature can be modeled either with an absorption edge ($E = 0.60_{0.33}^{0.75}$) or with an absorption trough ($E = 0.89_{0.80}^{0.94}$) with comparable probability ($\chi_r^2 = 0.72$ ($P = 0.72$) and $\chi_r^2 = 0.78$ ($P = 0.75$), respectively). However these features find no obvious interpretation at the redshift of the source. We then tried a warm absorber model (ABSORI of XSPEC), at the redshift of the quasar, obtaining a satisfactory fit ($\chi_r^2 = 0.89$). The fit requires a column density of the absorber equal to $N_H = 1.2_{1.2}^{10.3} \times 10^{22} \text{ cm}^{-2}$ and an ionization parameter $\xi = 262_{32}^{853}$, providing a reasonable physical interpretation. The results of the fits are reported in Table 4.

Previous ROSAT (Sambruna 1997) and ASCA (Kubo et al. 1998) spectra show a power-law X-ray spectrum with photon index $\Gamma_{2-10} = 1.7 \pm 0.1$, similar to that found with *BeppoSAX* . Notably these observations (with comparable flux) showed a smooth

continuum, without any evidence for spectral features.

The PDS/MECS normalization factor required by the data (2.7 ± 1.3) is well above the 90% upper limit (0.93) given by the SAX Cookbook. 0208-512 is located in a rich field: although the LECS and MECS images do not show other sources, a search using the NASA Extragalactic Database² reveals the presence of 13 sources catalogued as QSOs or X-ray sources, within a circular region with radius 1 deg centered on our target. It is then possible that the high-energy spectrum is contaminated by one or more sources present in the FOV (~ 1 deg) of the PDS.

In the fit presented here the PDS data are not included.

2.2. 0521-365

The LECS+MECS spectrum is well fitted by an absorbed power-law ($\Gamma = 1.77$ [1.73-1.85]) with column density consistent with the Galactic value. Previous X-ray observations with EXOSAT ($\Gamma \sim 1.6$, Pian et al. 1996) and ROSAT ($\Gamma = 1.92 \pm 0.05$, Sambruna 1997) are consistent with our results.

2.3. 1641+399 (3C345)

A power law model ($\Gamma \simeq 1.6$) with Galactic absorption fits the MECS+PDS data for $E > 2$ keV up to ~ 200 keV.

The inspection of the LECS image reveals the presence of several sources (not present in the MECS image): in particular a low-luminosity Seyfert galaxy, NGC 6212, located 5'

²<http://nedwww.ipac.caltech.edu/index.html>

away from the quasar hence within the standard extraction region of the LECS data. We conclude that the LECS spectrum could be contaminated by the presence of other sources close to 1641+399. In fact the residuals of a fit with an absorbed power-law model of the LECS data alone ($\Gamma \simeq 1.8$, $\chi_r^2 = 2.7$) show large scatter around the model (Fig.3), with the presence of possible features (with no straightforward interpretation) at energies ~ 2 keV and below ~ 1.0 keV. The features cannot simply be modeled as due to absorption of the continuum by a warm absorber at the redshift of the source.

A clear soft excess, interpreted as the hard tail of the UV bump, was present in the ROSAT data discussed by Sambruna (1997).

2.4. 2223-052 (3C446)

An absorbed power-law with Galactic absorption provides a good fit to the data. Our spectrum ($\Gamma \simeq 1.85$) is steeper than that measured on one occasion by GINGA ($\Gamma = 1.3$, Lawson et al. 1997). Moreover we do not confirm the ROSAT observations that seem to indicate an absorption column higher than the Galactic value at the 99% confidence level (Sambruna 1997).

2.5. 2243-123

A power-law with intermediate spectral index ($\Gamma = 1.7$) fits well the data.

The ROSAT observation reported by Siebert et al. (1998) showed a very steep spectrum, with $\Gamma = 2.9 \pm 0.5$ and a flux about 2.5 times larger than during our *BeppoSAX* pointing. Given the large flux variation, the steep spectrum measured by ROSAT could possibly represent intrinsic spectral variability possibly due to the high energy end of the

synchrotron component, extending up to the soft X-ray band.

2.6. 2251+158 (3C454.4)

As shown in Table 3 a fit with a power-law model and Galactic absorption gives an unacceptable fit to the data. Using a free value for N_H the fit improves significantly, requiring a column density higher than the Galactic value at the 99% confidence level (see the contour plot in Fig.4). The value of the required absorption column density in the quasar rest-frame is $N_{H,int} = 4.9_{2.4}^{7.9} \times 10^{21} \text{ cm}^{-2}$.

An acceptable fit ($\chi_r = 0.93$) of the low-energy portion of the spectrum can also be obtained with a broken power-law model (and Galactic column density), with a very flat ($\Gamma = -0.25$) soft spectrum. This, as in the case of other sources (e.g. 0836-710, Paper I), could be due to an intrinsic break in the continuum. From a statistical point of view both models have almost the same probability.

3. Modelling the SEDs

Using *BeppoSAX* and historical data we assembled the SEDs of the sources, shown in Fig.5. EGRET data are the average over the available observations as given by Hartman et al. (1999), while upper limits are from Thompson et al. (1995). For 0208-512 we report also the COMPTEL detection reported in Blom et al. (1995).

In the IR-optical bands, for 2223-052 are available quasi-simultaneous ISO (reported in Haas et al. 1998) and optical data. For other 3 sources (1641+399, 2243-123, 2251+158) we report average optical fluxes with the observed variability range. For 1641+399 Xie et al. (1999) report a variability amplitude in the *B*-band $\Delta B \simeq 3 \text{ mag}$ (which is shown in

Fig. 5), while for the other 2 sources the variability observed is rather small (less than 1 mag.). For 0208-512 and 0521-365 we report only the data we could find in literature. Both sources have IUE data. References to the data are reported in the figure caption.

The double-humped shape of the Spectral Energy Distribution (SEDs) is widely interpreted as due to synchrotron-Inverse Compton emission. The low energy component is synchrotron radiation from a population of relativistic electrons, while the high energy component is believed to be produced through Inverse Compton scattering of soft photons by the same electrons (although other scenarios have been considered, e.g. hadronic models, see Mannheim & Biermann 1992). The nature of soft photons involved in the IC process is still debated. In Blazars where thermal features (emission lines, the *Blue Bump*) are weak or even absent (as in BL Lac objects) the energy density of target photons is probably dominated by those produced by Synchrotron (SSC model, e.g. Maraschi, Ghisellini & Celotti 1992), while in powerful quasar (FSRQ) photons coming into the jet from the external environment form the dominant population (EC model, Dermer & Schlickeiser 1993; Sikora, Begelman & Rees 1994).

We have reproduced the overall SEDs using a Synchrotron-Inverse Compton (SSC+EC) model. The source is modeled as a sphere with radius R , tangled magnetic field with intensity B , in motion with bulk Lorentz factor Γ_b at an angle θ with respect to the line of sight. Relativistic effects are then regulated by the relativistic Doppler factor, given by: $\delta = [\Gamma_b(1 - \beta \cos \theta)]^{-1}$. Usually for Blazars $\theta \sim 1/\Gamma_b$, implying $\delta \simeq \Gamma_b$. Relativistic electrons emit through synchrotron and IC mechanisms. IC emissivity is calculated taking into account the full Klein-Nishina cross-section (following the Jones 1968 treatment). The electron energy distribution is modeled with the form:

$$N(\gamma) = K\gamma^{-n_1} \left(1 + \frac{\gamma}{\gamma_b}\right)^{n_1-n_2} \quad (1)$$

where K is a normalization factor, γ_b is the break Lorentz factor, n_1 and n_2 are the spectral

indices below and above the break, respectively. This particular form for the distribution function has been assumed on a purely phenomenological basis, in order to describe the curved shape of the SED.

The external radiation field is modelled as a black-body peaking at $\nu_{\text{ext}} \simeq 10^{15}$ Hz and energy density U_{ext} . The last quantity is calculated assuming that the radiation, with total luminosity τL_{ext} (τ , usually assumed to be ~ 0.1 is the fraction of the central emission reprocessed by the Broad Line Region) is diluted into a sphere of radius R_{BLR} . For 2251+158 which shows a clear UV bump we take $L_{\text{ext}} = L_{\text{UV}}$, while for the other cases we inferred L_{ext} from the luminosity of broad emission lines, following the approach used by Celotti et al. (1997). Luminosities are reported in Table 5. The value of R_{BLR} has been chosen to fit the observed data, but we checked that it is close to the value expected from the relation between the luminosity of the BLR and the radius derived (for radio-quiet quasars) by Kaspi et al. (2000). The external radiation energy density is amplified in the blob’s rest frame by a factor Γ^2 (e.g. Ghisellini et al. 1998). Moreover, as pointed out by Dermer (1995) (and confirmed by Georganopoulos, Kirk & Mastichiadis 2001), because of relativistic effects the beaming pattern of the EC radiation is narrower than that of the SSC emission. However, when $\delta \simeq \Gamma$ (as assumed here), the amplification is the same for both processes.

With these assumptions the model is well constrained by the observation and we can obtain fairly robust estimates of the principal physical quantities in the jet (see Paper I). The values of the parameters used to reproduce the SEDs are reported in Table 5. In all cases the bulk of the IC component is reproduced with the EC emission, while in some cases the SSC spectrum (peaking at lower energies) partly accounts for the low energy portion of the X-ray spectrum. The cut-off in the synchrotron spectrum around 10^{11} Hz is due to self-absorption. Clearly the adopted one-zone model can not account for the radio

emission, widely considered to be due to the superposition of the emission produced in the outer ($d \gtrsim 0.1$ pc) regions of the jet (e.g. Blandford & Konigl 1979).

4. Discussion and Conclusions

We have presented the results of the analysis of 6 FSRQs observed with *BeppoSAX*. The smooth power-law ($\Gamma = 1.6 - 1.7$) X-ray continuum is produced through Inverse Compton scattering on photon external to the jet, except for the “atypical” source 0521-365. In the case of two sources, 0208-512 and 2251+158, there is evidence of more complex X-ray spectra. In particular 0208-512 shows an absorption feature at ~ 0.6 keV, that we suggest could be explained allowing the presence of a warm absorption intercepting the emission produced in the jet.

The soft X-ray spectrum of 2251+158 shows a clear deficit of photons, consistent either with an intrinsic absorption (with rest-frame column density $\sim 5 \times 10^{21}$ cm $^{-2}$) or an intrinsic break in the continuum occurring below ~ 1 keV. In the latter case the position of the break could provide important insights into the physics of the jet, in particular regarding the value of the minimum energy of the radiating particles. On the other hand there is growing evidence that the presence of intrinsic absorption is common in high-redshift radio-loud quasars (e.g., Reeves & Turner 2000). Indeed *BeppoSAX* observations of two distant ($z \sim 4$) blazars (Fabian et al. 2001a,b) show the presence of huge absorption, with column density of the order of few 10^{23} cm $^{-2}$.

For three sources (0208-512, 1641+399 and 2251+158) we detect hard X-ray emission up to ~ 100 keV. For 1641+399 and 2251+158 the hard X-ray component appears to be the smooth extrapolation of the soft-medium X-ray continuum, while in 0208-512 the PDS spectrum is likely contaminated by another source present in the large FOV of

the instrument. We note however that 0208-512 is one of the few blazars detected by COMPTEL (Blom et al. 1995) and because of its exceptionally intense flux in the MeV band is considered as a prototype of the so-called “MeV Blazars”. Therefore we can not completely exclude the possibility that the spectral component responsible for the MeV flux contributes also in the PDS band, explaining the observed excess.

The modelling of the SEDs shows that FSRQs (except the case 0521-365, see below) are characterized by jets with Doppler beaming factors larger than 10 and magnetic field of the order of $B \sim 2 - 3$ G, roughly in equipartition with the emitting electrons. The size of the emitting region is of the order of 10^{16} cm, consistent with the absence of short-timescale variability ($\lesssim 1$ day) in this class of blazars. These results confirm, with a larger sample, the conclusions of Paper I

An interesting issue for Blazars is the value of the minimum Lorentz factor of the emitting electrons, $\gamma_{\min}$. In Paper I we showed how, from the absence of a spectral break in the soft X-ray continuum, it is possible to infer that the electron energy distribution extends down to $\gamma_{\min} \sim 1$. In fact a break in the EC continuum is expected at the frequency $\nu_{\text{br}} \simeq \Gamma^2 \nu_{\text{ext}} \gamma_{\min} \sim 10^{17} \Gamma_1 \sim$ (e.g. Sikora et al. 1997), corresponding to an energy ~ 1 keV (in the rest frame of the source). We recall that the lack of soft X-ray photons observed in 2251+158 (as for the case of 0836+710 reported in Paper I) could be interpreted as due to such intrinsic break in the continuum. Observations with larger signal to noise (possible with *XMM – Newton* or *Chandra*) will allow to disentangle the two possible interpretations, intrinsic curved continuum or absorption.

In our fits n_1 , the index of the low energy portion of the electron energy distribution, is smaller than 2. Clearly such a flat spectrum can not be produced by the cooling of high energy electrons (in that case one would expect $n_1 = 2$). On the other hand this spectrum is even flatter than the standard prediction $n_1 = 2$ for a population of non-thermal electrons

produced through Fermi acceleration by a non-relativistic shock (e.g. Blandford & Ostriker 1978). However recent calculations (see the review by Kirk & Dendy 2001) show that relativistic shocks, such as those likely present in the jet of blazars, could produce extremely flat ($n_1 < 2$) spectra.

The case of 0521-365 is atypical. Previous observational evidence suggested that the jet forms a relatively large angle with the line of sight. Pian et al. (1996), using independent arguments (in particular the presence of an optical jet, e.g. Scarpa et al. 1999), concluded that $\theta \sim 30^\circ$. Assuming that the bulk Lorentz factor of the emitting plasma is similar to that of the other blazars, $\Gamma_b \sim 10$, this implies that the emission from 0521-365 is weakly boosted ($\delta = 1 - 2$). The SED reported in Fig.5 has been calculated assuming $\Gamma_b = 10$ and $\theta = 15^\circ$, implying $\delta = 3$. In this situation the X-ray/ γ -ray continuum, contrary to the case of the other sources, is dominated by the SSC emission, while the EC spectrum gives only a small contribution in the γ -ray band.

Part of the motivation for this investigation was to study the difference between the EGRET and non-EGRET sources. In the whole FSRQ sample observed with *BeppoSAX* there are 9 EGRET sources and 3 sources not detected by EGRET. The average X-ray spectral indices ($\Gamma_{\text{EGRET}} = 1.56 \pm 0.06$ and $\Gamma_{\text{non-EGRET}} = 1.69 \pm 0.07$ for EGRET and non-EGRET sources, respectively) indicate that, within the statistical uncertainty, the X-ray spectral characteristics are similar for both groups. The modelling of the SEDS confirms the strong similarity between the two groups. The lack of the EGRET detection could be attributed to the variability behaviour typical of these sources in the γ -ray band (e.g. Mukherjee et al. 1997) rather than to an intrinsic difference in the properties of the jet. New γ -ray missions (AGILE, GLAST) will help to further investigate these problems.

We thank the *BeppoSAX* SDC for providing us with the cleaned data. This work was partly supported by the ASI grant I-R-105-27-00. This research has made use of

the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Bertsch, D.L. et al. 1993, ApJ, 405, L21
- Blandford, R. D. & Konigl, A. 1979, ApJ, 232, 34.
- Blandford, R. D. & Ostriker, J. P. 1978, ApJ, 221, L29
- Blom, J. J. et al. 1995, A&A, 298, L33
- Boella, G. et al. 1997, A&A, 122, 327
- Celotti, A., Padovani, P., & Ghisellini, G. 1997, MNRAS, 286, 415
- Comastri, A., Fossati, G., Ghisellini, G., & Molendi, S. 1997, ApJ, 480, 534
- Dermer, C.D. & Schlickeiser, R. 1993, ApJ, 416, 458
- Dondi, L., & Ghisellini, G., 1995, MNRAS, 273, 583
- Fabian, A. C., Celotti, A., Iwasawa, K., & Ghisellini, G. 2001a, MNRAS, 324, 628
- Fabian, A. C., Celotti, A., Iwasawa, K., McMahon, R. G., Carilli, C. L., Brandt, W. N.,
Ghisellini, G., & Hook, I. M. 2001b, MNRAS, 323, 373
- Falomo, R., Bersanelli, M., Bouchet, P., and Tanzi, E.G. 1993, AJ, 106, 11
- Fossati, G., et al. 1998, MNRAS, 299, 433
- Garcia, Aurea, Sodre', Laerte, Jablonski, Francisco J., Terlevich, Roberto, J. 1999, MNRAS,
309, 803
- Gear, W. K. et al. 1984, ApJ, 280, 102
- Geogory, P.C., Condon, J.J., 1991, ApJS, 75, 1011

- Georganopoulos M., Kirk J.G., Mastichiadis, A., 2001, accepted by ApJ
- Ghisellini, G., et al. 1998, MNRAS, 301, 451
- Ghisellini, G. et al. 1999, A&A, 348, 63
- Glass, I.S., 1981, MNRAS, 194, 795
- Green, P. J. 1996, ApJ, 467, 61
- Haas, M., Chini, R., Meisenheimer, K., Stickel, M., Lemke, D., Klaas, U., & Kreysa, E.
1998, ApJ, 503, L109
- Hartman, R. C., et al. 1999, ApJS, 123, 79
- Impey, C. D. & Neugebauer, G. 1988, AJ, 95, 307
- Impey, C. D. & Tapia, S. 1988, ApJ, 333, 666
- Jones, F.C., 1968, Phys. Rev., 167, 1159
- Kaspi, S., Smith, P. S., Netzer, H., Maoz, D., Jannuzi, B. T., & Givon, U. 2000, ApJ, 533,
631
- Kirk, J. G. & Dendy, R. O. 2001, Journal of Physics G Nuclear Physics, 27, 1589
- Kotilainen, T.K., Falomo, R., Scarpa, R. 1998, A&A, 336, 479
- Kubo, H., et al. 1998, ApJ, 504, 693
- Kuehr, H., Pauliny-Toth, I. I. K., Witzel, A., & Schmidt, J. 1981, AJ, 86, 854
- Lawson, A. J. & Turner, M. J. L. 1997, MNRAS, 288, 920
- Makino, F. 1989, Two Topics in X-Ray Astronomy, Volume 2: AGN and the X Ray
Background, 803

- Mannheim, K. & Biermann, P. L. 1992, A&A, 253, L21
- Maraschi, L., Ghisellini, G., & Celotti, A. 1992, ApJ, 397, L5
- Maraschi, L. et al. 1998, The Active X-ray Sky: Results from BeppoSAX and RXTE, 453
- Maraschi, L. & Tavecchio F., in “X-ray Astronomy 2000”, Proceedings to be published in ASP Conference series, eds. R. Giacconi, L. Stella, & S. Serio, 2000 (astro-ph/0107566)
- Moshir, M. E. A. 1990, IRAS Faint Source Catalogue, version 2.0 (1990)
- Mukherjee, R., et al. 1997, ApJ, 490, 116
- Neugebauer, G., Sofer, B.T., Matthews, K., Chanan, G.A., Margon, B., 1982, AJ, 87, 1639
- Padovani, P. & Urry, C. M. 1992, ApJ, 387, 449
- Patnaik, Alok R., Browne, Ian W. A., Wilkinson, Peter N., Wrobel, Joan M. 1992, MNRAS, 254, 655
- Pian, E. & Treves, A. 1993, ApJ, 416, 130
- Pian, E., Falomo, R., Ghisellini, G., Maraschi, L., Sambruna, R. M., Scarpa, R., & Treves, A. 1996, ApJ, 459, 169
- Pian, E. et al. 1999, ApJ, 521, 112
- Sambruna, R. M. 1997, ApJ, 487, 536
- Scarpa, R. & Falomo, R. 1997, A&A, 325, 109
- Scarpa, R., Urry, C. M., Falomo, R., & Treves, A. 1999, ApJ, 526, 643

- Siebert, J., Brinkmann, W., Drinkwater, M. J., Yuan, W., Francis, P. J., Peterson, B. A., & Webster, R. L. 1998, MNRAS, 301, 261
- Sikora, M. , Begelman, M. C. & Rees, M. J. 1994, ApJ, 421, 153
- Sikora, M., Madejski, G., Moderski, R., & Poutanen, J. 1997, ApJ, 484, 108
- Smith, P. S., Elston, R., Berriman, G., Allen, R. G., & Balonek, T. J. 1988, ApJ, 326, L39
- Stacy, J. G., Vestrand, W. T., Sreekumar, P., Bonnell, J., Kubo, H., & Hartman, R. C. 1996, A&AS, 120, 549
- Stevens, J. A., Litchfield, S. J., Robson, E. I., Hughes, D. H., Gear, W. K., Terasranta, H., Valtaoja, E., & Tornikoski, M. 1994, ApJ, 437, 91
- Tavecchio, F. et al. 2000, ApJ, 543, 535 (Paper I)
- Thompson, D.J., et al. 1995, ApJS, 101, 259
- Tornikoski, M., et al. 1996, A&AS, 116, 157
- Urry, C.M. and Padovani, P. 1995, PASP, 107, 803
- Villata et al. 1997, A&AS, 121, 119
- Wiren, S., Valtaoja, E., Terasranta, H., Kotilainen, J. 1992, AJ, 104, 1009
- Wright, A. E., Ables, J. G., Allen, D. A. 1983, MNRAS, 205, 793
- Wright, A. & Otrupcek, R. 1990, PKS Catalog (1990)
- Wright, Alan E., Griffith, Mark R., Hunt, Andrew J., Troup, Euan, Burke, B.F., Ekers, R.D. 1996, ApJS, 103, 145
- Xie, G.Z., Zhang, X., Bai, J.M., and Xie, Z.H. 1998, ApJ, 508, 180

Figure Captions

Fig. 1.— Fit of LECS and MECS data of 0208-512 with a power-law and free absorption. The feature at ~ 0.6 keV is clearly visible in the data to model ratio (lower panel).

Fig. 2.— Fit of LECS and MECS data of 0208-512 with a power-law (Galactic absorption) plus an absorption edge.

Fig. 3.— Fit of LECS data of 1641+399 with a power-law and Galactic absorption.

Fig. 4.— 68, 90 and 99% photon index- N_H confidence contours for 2251+158. The column density exceeds the Galactic value at >99% level.

Fig. 5.— Spectral Energy Distributions of the sources analyzed in this work. The solid line is the model discussed in the text. References for the data: **0208-512**: Bertsch et al. 1993, Tornikoski et al. 1996, Stacy et al. 96, Impey & Tapia 1988, Blom et al. 1995 **0521-365**: Wright & Otrupcek 1990, Wright et al. 1996, Glass et al. 1981, Falomo et al. 1993, Xie et al. 1998. **1641+339**: Wiren et al. 1992, Patnaik et al. 1992, Geogory & Condon 1991, Moshir et al. 1990, Neugebauer et al. 1982, Villata et al. 1997, Xie et al. 1998. **2223-052**: Wright & Otreepcek 1990, Haas et al. 1998, OPTICAL. **2243-123**: Wright & Otreepcek 1990, Wright et al. 1983, Kotilainen et al. 1998, Garcia et al. 1999. **2251+158**: Kuhr et al. 1981, NED, Gear et al. 1994, Stevens et al 1994, Impey & Neugebauer 1988, Smith et al. 1988, Pian & Treves 1993.

Table 1: List of the sources discussed in the present work. The last column indicates EGRET detection or lack thereof.

Source	Other name	z	EGRET
0208-512	PKS	1.003	Y
0521-365	PKS	0.055	Y
1641+399	3C 345	0.593	N
2223-052	3C 446	1.4	N
2243-123	PKS	0.63	N
2251+158	3C 454.4	0.859	Y

Table 2: *Beppo*SAX data observation log.

Date	Start	End	LECS Exp.(s)	net cts/s ^a	MECS Exp.(s)	net cts/s ^b	PDS Exp.(s)	net cts/s
0208-512								
14-15/01/01	13:37:10	13:00:09	15650	0.022 ± 0.001	34256	0.053 ± 0.001	15260	0.1417 ± 0.03
0521-365								
2-3/10/98	10:16:24	09:46:23	17752	0.067 ± 0.002	41082	0.106 ± 0.001	19052	0.05 ± 0.04
1641+399								
19/02/99	03:55:43	20:13:57	10934	0.045 ± 0.002	25866	0.059 ± 0.002	13052	0.15 ± 0.05
2223-052								
10-11/11/97	13:22:27	00:22:40	9200	0.01 ± 0.001	16180	0.016 ± 0.001	6860	0.008 ± 0.08
2243-123								
18-19/11/98	18:41:31	11:49:58	10060	0.015 ± 0.002	27500	0.022 ± 0.001	13222	0.004 ± 0.015
2251+158								
5-6/6/2000	15:36:05	21:44:38	17425	0.046 ± 0.001	48515	0.115 ± 0.002	22439	0.29 ± 0.03

^a: 0.1-4 keV

^b: 1.8-10.5 keV, 2 MECS units

Table 3: Absorbed power-law fits to *Beppo*SAX data. Errors are quoted at the 90% confidence level for 1 parameter of interest.

Source	Γ^a	N_H	$F_{[2-10\text{ keV}]}$	$\chi^2/\text{d.o.f.}$
		10^{20} cm^{-2}	$10^{-12} \text{ erg cm}^{-2} \text{ s}^{-1}$	
0208-512 (LE+ME2+PDS) ^b	$1.68^{1.80}_{1.51}$	$16.7^{30.0}_{0.1}$	4.7	51.16/28
0521-365 (LE+ME)	$1.77^{1.85}_{1.73}$	$4.3^{7.6}_{3.2}$	8.72	69.35/89
1641+399 (LE+ME+PDS)	$1.57^{1.66}_{1.47}$	1.12(fix)	5.2	65.26/45
2223-052 (LE+ME)	$1.84^{2.27}_{1.56}$	$7.1^{76.1}_0$	1.25	14./16
-	$1.86^{2.15}_{1.58}$	4.87 (fix)	-	14.72/17
2243-123 (LE+ME)	$1.73^{1.91}_{1.56}$	$3.05^{7.7}_{1.28}$	1.83	18.37/23
2251+158 (LE+ME+PDS)	$1.31^{1.26}_{1.36}$	6.5(fix)	10.6	94.2/80
-	$1.38^{1.33}_{1.44}$	$18.6^{25.1}_{12.2}$	-	77.1/79

^a: photon index, related to the spectral index by $\alpha = \Gamma + 1$.

^b: see Table 4

Table 4: Fits to *BeppoSAX* data of 0208-512. Errors are quoted at the 90% confidence level for 1 parameter of interest.

Model	Γ^a	E_{abs}/N_H^b	τ/ξ^c	$\chi^2/\text{d.o.f.}$
PL+edge	$1.72_{1.65}^{1.88}$	$0.56_{0.33}^{0.75}$	$2.1_{0.9}^{10}$	13.25/17
PL+notch	$1.68_{1.55}^{1.81}$	$0.89_{0.80}^{0.94}$	$0.21_{0.13}^{0.31}$	12.85/17
PL+ABSORI	$1.70_{1.62}^{1.96}$	$4.2_{1.2}^{10.3}$	262_{32}^{853}	15.13/17

^a: photon index, related to the spectral index by $\alpha = \Gamma + 1$.

^b: Energy of the absorption feature (keV) or N_H for ABSORI (in units of 10^{20} cm^{-2}).


^c: optical depth for the edge, width (keV) for NOTCH or ξ for ABSORI

Table 5: Parameters used for the emission model described in the text.


R	B	δ	γ_b	n_1	n_2	K	L_{BLR}	R_{BLR}
10^{16} cm	G					cm^{-3}	10^{45} erg s $^{-1}$	10^{18} cm
0208-512								
1.5	1.5	18	100	1.4	3.8	2×10^4	2.1	0.6
0521-365								
2.0	0.3	3*	8.8×10^3	1.25	4	3×10^3	2.4×10^{-2}	0.8
1641+399								
4.0	2.9	9.75	200	1.5	4.2	2.8×10^3	3.7	0.6
2223-052								
4.25	5.6	17	135	1.6	4.3	1.7×10^3	8.5	1.1
2243-123								
3.5	2.5	15	250	1.6	4.3	1.7×10^3	5.6	0.9
2251+158								
4.0	1.5	12	60	1.8	3.4	5×10^4	4.	1.0

*: see text


0208–512 – Power law



0208–512 – PL+Edge



1641+339 – LECS – PL



2251+158 – Absorbed power-law

